

表四、抗番茄捲葉病毒病番茄新品系區域試驗園藝特性調查

Table 4. Horticultural characteristics of tomato hybrid lines (F₁) at regional trial.

Crop	Entry	Average fruit weight (g)	Fruit setting (%)	Incidence rate of ToLCV (%)	Soluble solid (°Brix)	Acidity (%)	Fruit thickness (mm)	Immature fruit shoulder color
2003 summer/ autumn	FMTT957	137.6 ^{b1}	54.3 ^a	6.7 ^b	5.7 ^a	0.42 ^b	6.9 ^a	Dark green
	FMTT965	142.6 ^b	52.9 ^a	6.0 ^b	5.2 ^a	0.48 ^a	6.4 ^a	Green
	Taichung Asveg 10 (CK)	168.3 ^a	53.6 ^a	36.5 ^a	5.6 ^a	0.40 ^b	6.4 ^a	Dark green
2004 Spring	FMTT957	146.5 ^b	55.5 ^a	8.8 ^b	5.3 ^a	0.46 ^a	5.4 ^a	Dark green
	FMTT965	141.6 ^b	59.7 ^a	7.4 ^b	5.3 ^a	0.43 ^a	5.3 ^a	Green
	Taichung Asveg 10 (CK)	172.7 ^a	57.4 ^a	82.7 ^a	4.7 ^b	0.44 ^a	4.8 ^b	Dark green

¹ Means followed by the same letters are not significantly different at 5% level.